

Edex—a reliable choice.



Local Adverse Reactions Reported by ≥5% of Patients (n=1065) All Study Periods*

Local Reaction	Edex® N = 1065 n (%)
Penile pain during injection	305 (29)
Penile pain during erection	368 (35)
Penile pain after erection	317 (30)
Penile pain (other)**	116 (11)
Bleeding	158 (15)
Hematoma	56 (5)
Penile angulation	72 (7)
Penile fibrosis	52 (5)
Faulty injection technique***	59 (6)

* Protocol Numbers KU-620-001, KU-620-002, KU-620-003, F-8653.

** Penile pain reported without an association to injection site or erection, such as pain in penis and scrotum, pain in glans penis, and burning penile pain.

*** Examples include injection into glans penis, urethra or subcutaneously.

Systemic Adverse Events Reported by ≥5% of Patients in Clinical Studies (n=1065)*

- Upper respiratory tract infection n=58/1065 (5%)

Please see full Prescribing Information.


1. Montorsi F, Salonia A, Deho F, et al. Pharmacological management of erectile dysfunction. BJU Int. 2003;91(5):446-54.

Count on Edex

- Only dual chamber injection (alprostadil) system currently available
- No refrigeration required

Prescribe with confidence

- FDA approved and on the market for over 10 years
- Financial support for qualified patients



Because oral ED* therapies are not for everyone. 1

edex®
(alprostadil for injection)
10 mcg • 20 mcg • 40 mcg
➤ A Reliable Choice.

actient
pharmaceuticals™

Actient is a new specialty pharmaceutical company dedicated to serving the urology community with therapies like Edex. Our team of sales professionals have extensive experience representing urology products and will be supporting you with Edex information and patient support.

© 2010 Actient Pharmaceuticals, Inc., Deerfield, IL 60015 USA ACT-M1001 09/10 All rights reserved.
Edex® is a registered trademark of UCB group of companies.

edex®
(alprostadil for injection)
10 mcg • 20 mcg • 40 mcg

*Erectile Dysfunction


Prescribe Edex—a reliable choice.



For men who don't respond to or aren't candidates for oral therapy.¹

Start with reliable efficacy

The mean rate of response with an erection sufficient for intercourse*



* Patients received in-office evaluations, dose titration, and proper training techniques prior to the 2-week placebo-controlled phase which was followed by the open-label, at-home extension period, which ranged from 6 to 12 months. Maximum allowable dose was 40 mcg.

Edex is indicated for the treatment of erectile dysfunction due to neurogenic, vasculogenic, psychogenic, or mixed etiology.

Edex® is not a cure for erectile dysfunction. The underlying treatable medical causes should be diagnosed and treated prior to initiation of therapy. The therapeutic effect of each dose is temporary. Edex® should be used no more than 3 times per week with at least 24 hours between each dose.

If priapism occurs, the patient should seek immediate medical attention.

Count on Edex the only dual chamber (alprostadil) injection system currently available

- Three strengths: 10, 20 and 40 mcg
- Supplied in 2- and 6-cartridge packages
- No refrigeration required

Prescribe Edex with confidence

- FDA approved
- Greater than 10 years of patient experience
- Rapid onset of action (5-20 minutes)
- Edex should be titrated slowly to the lowest effective dose
- To ensure safe and effective use of Edex, the patient should be thoroughly instructed and trained in the self-injection technique before he begins intracavernous treatment with Edex at home

Edex should not be used in men for whom sexual activity is inadvisable or contraindicated.

Edex should not be used in women and children and is not for use in newborns.

Edex® is contraindicated in men with known hypersensitivity to alprostadil or other prostaglandins, men with conditions that might predispose them to priapism, and patients with penile implants or anatomical deformities of the penis.

Please see back page for additional safety information.



Financial support for qualified patients

To the Patient: You must present this card to the pharmacist along with your prescription to participate in this program. If you have any questions regarding your eligibility or benefits, or if you wish to discontinue your participation, call the Edex Loyalty Card program at 1-800-657-7613 (8:00 AM-8:00 PM EST, Monday-Friday). When you use this card, you are certifying that you understand the program rules, regulations, and terms and conditions. You are not eligible if prescriptions are paid by any state or other federally funded programs, including, but not limited to Medicare or Medicaid, Medigap, VA or DOD or TriCare, or where prohibited by law; and you will otherwise comply with the terms above. Only male patients who are 18 years or older can participate in this program. Each card provides one benefit per patient for the life of the program.

To The Pharmacist: When you use this card, you are certifying that you have not submitted and will not submit a claim for reimbursement under any federal, state or other governmental programs for this prescription.

- Submit transaction to McKesson Corporation using BIN #610524
- If primary coverage exists, input card information as secondary coverage and transmit using the COB segment of the NCPDP transaction. Applicable discounts will be displayed in the transaction response.
- Acceptance of this card and your submission of claims for the Edex Loyalty Card program are subject to the LoyaltyScript® program Terms and Conditions posted at www.mckesson.com/mprstnc
- Patient is not eligible if prescriptions are paid in part or full by any state or federally funded programs, including but not limited to Medicare or Medicaid, Medigap, VA, DOD or TriCare and where prohibited by law.
- Cash payers are eligible for this program.
- For questions regarding setup, claim transmission, patient eligibility or other issues, call the Edex Loyalty Card program at 1-800-657-7613 (8:00 AM-8:00 PM EST, Monday-Friday).**

edex®
(alprostadil for injection)
10 mcg • 20 mcg • 40 mcg
A Reliable Choice.